

RESOURCE FOR DANCERS AND TEACHERS

Nutrition Resource Paper

JASMINE CHALLIS RD, AND ADRIENNE STEVENS EDD, WITH THE IADMS DANCE EDUCATORS' COMMITTEE, 2016, AND UPDATED IN 2019

CONTENTS:

1. INTRODUCTION
2. FOOD AND DIETARY RECOMMENDATIONS
3. FLUID, HYDRATION, AND SWEAT
4. RECOMMENDED READING
5. REFERENCES
6. AUTHORS

1. INTRODUCTION

To perform at their best, dancers need to be adequately fueled for the activities in which they participate regularly: classes, rehearsals, and performances/competitions. This paper will present concise and practical strategies regarding the types and amounts of food that are needed to sustain health across the variety of activities dancers participate in, specifically addressing a balance of nutrients: carbohydrates, fats, proteins, micronutrients and fluids. In addition, it will stress the practical assessment of food in context to the amount of energy exerted in relation to rest and recovery rather than focusing on calories alone. With this in mind, we propose that dancers develop greater self-awareness of these important components when considering food choices and timing: amount of time spent dancing, intensity, type of dance activity, individual metabolic variation, access to food and cooking, and personal goals.

Adequate nutrition is intricately tied to every aspect of physiology and health. To that point, weight management is often on many dancers' minds and can end up dominating thinking in an unhelpful way. Being clinically underweight or overweight can trigger emotional issues or bone, hormonal, and metabolic abnormalities that can last a lifetime.¹ In contrast, in addressing this approach, this paper stresses the importance of health and fitness recognizing how nutrition fits into dancers' lives. Because each dancer is different, individualized approaches to diet, goals, and dance aesthetic must be taken into consideration.

GETTING THE FACTS

As a population, dancers often obtain eating and dietary advice from various sources that are not always credible, and instead take recommendations made by celebrities, the media, and from word of mouth. Often, nutritional advice by qualified healthcare professionals is not sought out.² Wherever possible, food and eating recommendations should be obtained from credible sources based on current medical and scientific data and this information should be evaluated by the dancer. Dietitians (or notably, Sports Dietitians) are the most specifically qualified experts in this field. Diets that promote rapid

weight loss and typically exclude major food groups or require the addition of herbs and supplements, are likely to impair performance, are not sustainable, can be harmful to health, and should not be followed. In addition, self-proclaimed experts who advertise commercial products hyped to produce large and rapid gains in muscle mass, stimulate metabolism, or provide energy, should be considered with skepticism.

Some of the characteristics that define dance training include the ability to apply intelligence, discipline, self-discovery, and focus. In striving for excellence however, dancers often take an 'all or nothing' approach, thinking that if rules are obeyed to the letter they will be better performers, have better technical skills, be thinner, healthier, or be better individuals (insert any superlative here). Dancers are bombarded with media information and jargon that may not pertain to their needs. Unfortunately, dancers often erroneously believe that if one type of food, vitamin, or exercise is good for them, much more is better. ² This approach can result in nutritional imbalance and is not sustainable. While it would be easier if nutrition advice for the dancer could be presented in black and white terms, the reality is that nutrition is not like this. Eating behavior is driven by complex neuro-hormonal mechanisms which regulate cravings, appetite and satiety and physiological responses to food. Dancers, like the general population, are hard-wired for salty, sweet, and high fat foods, yet there is limited room for these foods in a dancer's diet. It is the combination and variety of foods which matter, and this takes work to plan. For example, a diet made up of large amounts of vegetables and whole grains will not result in optimal performance for most dancers, nor will one which is based only on candy and cake. We stress that consuming a variety of foods is necessary for the body and the mind and to remember that although food is fuel, eating should be pleasurable. Finally, global food supply and local food issues should be researched by the dancer. Opting to eat vegan, vegetarian, or engage in other ethical decisions around food should be choices that are informed by both circumstance and research and this is beyond the scope of this paper. For information on plant based diets, see: <http://www.mayoclinic.org/healthy-lifestyle/nutrition-and-healthy-eating/in-depth/vegetarian-diet/art-20046446>

The dancer must develop a performance eating plan to navigate challenges and optimize their condition in order for their body and mind to function at a high level. When the body is given what it needs to function at a high level, the need for overly restrictive eating is minimized. Since every person is different, when possible, seek out nutritional advice from a qualified nutrition professional (such as a Sports Dietitian). As qualifications and licensure requirements of those providing nutritional counseling is varied and differs by country and state, research may be needed.

For information on nutrition credentialing

https://www.bda.uk.com/publications/dietitian_nutritionist.pdf

<http://www.nutritionaustralia.org/national/nutritionist-or-dietitian-which-me>

DANCERS AS PERFORMING ATHLETES

Dancers are artistic athletes. The performances they undertake can be lengthy and physically demanding. They need to consume sufficient quantities of energy (E), or fuel, from the major food groups to preserve metabolic function, hormonal and growth regulation, as well as to meet the demands required by their activity. This can be difficult to manage since the dance aesthetic is generally leaner for both men and women compared to non-dancer standards. For example, body size is frequently defined relative to body mass index (BMI). There are many web-based tools that can calculate one's BMI based on knowing and inputting one's height and weight. BMI is a ratio of height to weight, with the normal range for adults between 20-25 kg/m². This range represents the optimum for long term health as derived from large scale population health data suggesting that adults within this range carry a lower risk of chronic and debilitating diseases (cardiovascular disease, diabetes, cancer, etc.) compared to those who are overweight, obese, or underweight.³ A caveat of BMI, and particularly for the dancer who has proportionally more muscle to fat, is that this measurement does not reflect body composition. Therefore, a dancer's weight may be high as a result of muscle, when, in reality, the proportion of weight attributable to fat is low. In non-dancer/athletes, normal adults with a BMI below 17 usually translates to lower body fat and body muscle which can impact long term health, whereas a BMI over 30 in most cases indicates excess body fat which negatively impacts health. Since BMI scores are calculated differently for adults

and children, dancers below the age of 18 should consult growth charts, or a medical professional who can make an accurate assessment of growth and body size, to verify age-related ranges for BMI.

To calculate BMI: https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi-m.htm

Link to growth charts: <http://www.infantchart.com/child/>

DIETING VERSUS A HEALTHY BODY WEIGHT

There are intrinsic and extrinsic variables in maintaining a healthy body weight. Dancers are bombarded by conflicting information that they must sort through. A dancer cannot control expectations of the dance aesthetic or cultural views about ideal body types. Body type is individual, and body composition is primarily determined by one's genetic profile. Dancers should be encouraged to look at the frame sizes of close family to see how to best work with their body, rather than against it, in order to have a fit, healthy and sustainable dancer's body. Maintaining health and the ability to function well is most critical. Dancers can control their food intake and calculate their energy expenditure. Dancers can control the way that sleep affects their performance. Sleep is an important component of appetite regulation, and lack of sleep can make it difficult to avoid excessive hunger as the body looks for energy.

There are many diets, supplements and tools that are promoted by celebrities, pop artists, athletes, and even famous doctors in social and traditional media that boast exaggerated claims to enhance performance, promote muscle mass, decrease body weight, take away hunger, etc. Many of these claims are not validated by scientific research in humans or regulated for safety and purity by government drug agencies, and can range from mildly effective to ineffective, dangerous, and/or a waste of hard-earned money. There is no magic pill to produce quick health results and promises that sound too good to be true are usually not. Any changes to body composition will need to be made over weeks to months, not in a few days.

DISORDERED EATING

Unfortunately, disordered eating and eating disorders are common in dancers. Dancers are advised to consult a specialist for a personalized plan or program in the attempt to lose or gain weight. Disordered eating can be described as any eating or food-related experiences or behaviors that impacts negatively on a person's health or wellbeing. Engaging in disordered eating behavior is not necessarily diagnostic of having an eating disorder, but there can be many similarities or overlapping factors at play.

Some examples of disordered eating are:

- Applying all-or-nothing "rules" around eating that are stressful or difficult to manage or maintain over time
- Avoiding eating with others
- Going for long periods of time without eating or deliberately skipping meals
- Counting calories, or grams of food in a way that is unhelpful, stressful, or extremely regimented
- Feeling guilt or shame around food or eating
- Feeling anxious around food or eating

The most defining characteristic of eating disorders are the eating behaviors, rather than body size. Seeking support or help for an eating disorder from qualified and experienced professionals is absolutely critical, particularly for dancers who are looking to continue dancing and take care of their bodies in a more positive and healthy way.⁴

If you suspect a friend of having an eating disorder, don't stay silent! Tell a teacher, parent, or someone with authority that can help. You can save a life.

For more information please see National Eating Disorder websites:

UK: <http://www.b-eat.co.uk>

US: <https://www.nationaleatingdisorders.org/>

AUS: www.nedc.com.au/

THE DANCE AESTHETIC

Each type of dancing has its own body ideal, with female ballet dancers typically having the lowest body weight compared to most other dance forms. A vast amount of literature, anecdotal reports, and portrayals in books and film illustrates the risk that dancers face in pursuing thinness, resulting in disordered eating. Studies in dancers, models, and athletes have shown a higher prevalence of eating disorders and poor body image compared to individuals who do not partake in these activities. Recently, more information has emerged documenting eating disorders in men.¹ Thus the prevention of eating disorders should be equally targeted at men and women. Because a dancer relies on their body, optimal physical training balanced with good nutritional practice allows the dancer to train and perform at their best. Maintaining health by achieving this balance is essential both in the short term and all throughout the dancer's career. Fortunately, this is a concern in the dance world as well. For example, prestigious competitions such as the Prix de Lausanne have set minimum safe weights that dancers must be *above* in order to participate.⁵

FINDING THE RIGHT BALANCE

A number of factors should be considered in the recommendation of a nutritionally-balanced plan for dancers including type of training frequency and intensity, timing of eating in relation to activity, health and body weight goals. Although dancers spend much time perfecting their craft and exerting energy in the studio, stage or in rehearsal, in actuality, the amount of energy expenditure (EE) and metabolic demand is frequently lower than they assume.⁶ For example, the demands of a ballet class which requires extreme concentration, effort, and endurance in a standing position at the barre or during center work, frequently are aerobically low. These combined elements can cause muscle soreness and fatigue, and while perceived as major exertion, metabolic demand is low and very few calories are burned in the process. The intensity and frequency of classes will determine the dancer's nutritional requirements. Some dance styles, however, such as Irish, tap or jazz, do include high intensity long routines which may be more energy-

demanding. Therefore the basic tenets of physical activity assessment begin with awareness of quantitative and qualitative factors. While there is individual variability around an average, to evaluate the type of metabolic demands based on activity, dancers should assess their frequency, intensity, timing, and the type of training (FITT), a common exercise science assessment. Beyond this, however, we know that the body has a requirement for maintaining physical and brain function.^{7, 8}

Aerobic activity improves or is intended to improve the efficiency of the body's ability to produce energy through the utilization of carbohydrate and free fatty acids (FFA) as its main fuel source. An increase in aerobic capacity allows a greater utilization of FFA during low to moderate dance activity thereby sparing carbohydrate for high intensity activities. These systemic responses protect the body against more serious disease, build endurance, enhance glucose metabolism, help maintain normal weight and improve sense of well-being.

Anaerobic activity is high intensity and is used in short duration exercise bursts. This type of training often produces lactate resulting in fatigue. For more information, see the IADMS Resource paper on Dance Fitness: <https://www.iadms.org/?303>

Determining energy expenditure (A FITT SELF-Test)

Frequency: How many classes/rehearsals/performances/supplemental training bouts/per day?

Intensity: How hard am I physically working? Does the activity increase heart rate and cause rapid breathing? Examples: Petite allegro, jumps across the floor, fast turning combined with leaps and covering a large amount of space.

Does the activity require sustained muscular strength without much movement?
Examples: Most barre work, center adagio.

Timing: How much time (minutes/hours) is spent dancing?
How long are the activities sustained? For dancers, this can vary by the type of class.

Training: Different types of dancing require different energy needs.

Different styles require either whole body activation or just fancy footwork. In general, the more body parts are involved combined with accelerated breathing rates, the more aerobically demanding.

Examples: Rapid tap dancing or break dancing compared to a Sarabande or slow waltz.


FEMALE ATHLETE TRIAD AND RELATIVE ENERGY DEFICIENCY IN SPORT

The Female Athlete Triad (“The Triad”) and Relative Energy Deficiency in Sport (RED-S) are of great concern in sports but also are important topics for dancers and dance educators. The Triad is a medical condition first described over 30 years ago, seen in physically active girls and women characterized by low energy availability with or without disordered eating, menstrual dysfunction, and low bone mineral density. When one or more of the three Triad components is detected, early intervention is essential to prevent its progression.⁹ RED-S refers to impaired physiological and psychological function caused by low energy availability (LEA). LEA, which underpins the concept of RED-S, is a mismatch between a dancer’s energy intake (diet) and the energy expended in exercise, leaving inadequate energy to support the functions required by the body to maintain optimal health and performance. RED-S includes the menstrual dysfunction and low bone mineral density affecting only women identified in The Triad, together with the impact of low energy availability on many other body systems, including the digestive system, the immune system, the musculoskeletal systems and the hormonal systems all of which can affect both men and women. The consequence is that this energy imbalance affects health, growth, repair as well as daily living and sporting activities.¹⁰ There is acceptance that more research is needed in this field, but the 2018 update is very clear that the consequences of low energy availability can be severe and that this is a condition which needs careful management to prevent long term consequences.

2. FOOD AND DIETARY RECOMMENDATIONS FOR DANCERS

The recommendations below offer basic guidelines which have been adapted from sports nutrition research. A specialist should be consulted for a personalized nutrition

prescription. Remember that a dancer's needs are unique and sometimes nutritional choices are influenced by medical conditions. In general, the energy in a dancer's diet should be composed of about 55%-60% carbohydrates (CHO), 12%-15% proteins (P) and 20%-30% fats (F). CHO, F, and P are necessary components the human body needs to maintain normal physiologic function. All dancers need to ingest sufficient energy to meet the rigors of training. Consuming the right amounts and types of food and fluid will provide the body with the "high performance fuel" necessary to achieve optimal training benefits and peak performance. Because every person is different, many factors including food intolerances, allergies, cultural and religious reasons affecting food choice must be taken into consideration when devising any dietary program. Not only is what a dancer eats important, but when and how much needs to be critically evaluated as well.



Recommendations for dancers vary slightly compared to non-athletic adults whose intake would be more varied: 45%–65% CHO, 20%–35% F, and 10%–35% P. Protein is recommended in absolute amounts. The recommended dietary allowance (RDA) for protein is 46g/day for women and 56 g/day for men or 0.8 g/kg body weight of P (1kg=2.2lb).¹¹ Female dancers are advised to aim for 1g protein per kg body weight, while male dancers should aim for 1.5g protein per kg body weight and up to 2g per kg body weight if aiming to increase muscle mass. For more information on carbohydrates,

proteins and fats scroll down through the paper. Each section contains reference charts with recommended amounts for typical foods in the group.

In addition, reading food labels or using nutrition apps can help roughly calculate the day's nutrients. Limiting or restricting any macronutrient [fats, proteins or carbohydrates] group decreases feelings of fullness (satiety), both during the meal and thereafter. A combination of each nutrient in the right proportion is best in order for the correct signaling between the body and brain to occur. Each nutrient has a unique contribution to the body so over- or under- consuming one or another nutrient is not beneficial. On the other hand, under consumption of food is detrimental to bone health as well as energy levels.

Use this link for information on variety of proteins from various food sources:

<http://www.choosemyplate.gov/protein-foods>

Carbohydrates: How much do you need?

<https://www.sportaus.gov.au/ais/nutrition>

CARBODYDRATES

Glucose (simple sugar) is the sole fuel for the brain and a major fuel, together with glycogen, in the muscles. CHO are broken down into glucose in the digestive tract to be used to maintain blood sugar levels, to fuel the brain and are also stored in muscle (350g to 400g) and in the liver (100g) in the form of glycogen. However, the body storage of glucose (glycogen) is limited and sugars are converted to fat when the reserves are filled. CHO supply is critical for strength and endurance.

Not all CHOs and sugars have equivalent nutritional value. Dancers tend to think that all CHO and sugars are inherently bad, are the source of empty calories (i.e. calories with no vitamins, minerals, or protein) and cause weight gain, but this is not true. Many foods contain CHO, in the form of starches and sugars found naturally within the food. Those foods where the CHO (whether sugar or starch) is more processed, for example, white

bread, white rice, cookies/biscuits, fruit juice, sweets, sweet drinks, and chocolate, are generally less helpful to dancers than those which are less processed. The more the food is processed, the fewer the micronutrients it contains. More processed foods contain fewer micronutrients and are easier to over consume. Nutrient dense foods sustain energy and physiologic processes better over time. Dancers are advised to be careful about consuming too few CHO when they are in rigorous training or performance as insufficient CHO consumption compromises the ability to sustain energy which contributes to fatigue. As a guide dancers will typically need between 4 and 8 grams of carbohydrate per kg body weight. This will vary according to the duration and intensity of the workload, and at times may be even higher (during long or intense rehearsals or demanding pieces).¹²

TABLE I: FOODS RICH IN CARBOHYDRATES

TABLE 1: FOODS RICH IN CARBOHYDRATES				
Food	Type of CHO	Benefits	Recommendations	Portion size
Bread, breakfast cereals including oatmeal, instant oatmeal or porridge, muesli and granola, rice	Combination of slowly digestible and rapidly digestible starch with small amounts of natural sugars (plus added sugars in most cereals).	Whole grain types contain more vitamins, minerals and fiber than white versions and generally digest more slowly, helping keep blood glucose levels stable. Finely ground oats are digested faster.	Muesli and granola can vary in sugar content. Best to choose one without added sugar.	75g-125g bread 45g-75g cereal/rice
Pasta, including whole wheat and those with additions such as spinach or tomato	Higher proportion of slowly digestible starch than foods in box above.	Whole grain pasta has a higher amount of fiber and B vitamins. All pasta is digested more slowly than many other CHO rich foods.	A good choice; check portion sizes.	45g-75g dry weight
White potatoes, sweet potatoes	Combination of slowly digestible	New/baby potatoes and	Avoid chips, french fries, or	180g-300g

	and rapidly digestible starch with small amounts of natural sugars.	sweet potatoes are best for slow release of energy.	roasted potatoes as they can be high in fat.	
Root vegetables including parsnips, carrots, turnips	All have much less carbohydrate than potato. Parsnip and beetroot have 50% CHO as found in potato.	These supply a range of vitamins and chemicals helpful to long-term health.	Include as wide a range of vegetables as possible.	60g+
Beans and lentils	Slowly digestible starches, negligible sugars.	Good source of protein also. Provides a number of vitamins and minerals including iron.	Include regularly if digestion allows. If using for both protein and CHO, large portions will be needed.	60g dry/190g cooked- 100g dry/320g cooked
Quinoa	Combination of slowly digestible and rapidly digestible starch with small amounts of natural sugars.	Slightly higher content of protein compared to with many grains. The slow release of CHO is to sustain energy levels over time.	Often expensive so use as part of a variety of CHO.	45g-75g dry weight
Fruit	Combination of glucose, fructose and sucrose.	2-3 whole fruit per day supply natural CHO, fiber, vitamins and minerals.	Whole fruits provide better alternatives to fruit juices, which should be limited.	1 whole fruit approx. 120g
Milk, yogurt including Greek yogurt and Greek style yogurt	Natural milk sugar, lactose, sweetened yogurt have added sucrose.	Good source of calcium and protein.	Choose unsweetened yogurt and add fruit and nuts for best nutrition.	200ml milk (1 medium glass) 150-180g yogurt (1 cup)
Cakes, cookies/biscuits, sweets, chocolate, sweetened drinks, both carbonated and still/non-carbonated, liquid yogurt drinks	Starch/Sugars, of which most is added refined sucrose.	Likely to result in rapid increases in blood glucose levels which will not be sustained and can contribute to acne.	They lack other nutrients and are poor choices to maintain energy needed for dancing.	

FAT

Why do we need fat? A mix of fat and glucose is needed for energy during exercise and at rest. Dietary fat is essential for the regulation of multiple physiologic systems; it is needed for the absorption of fat soluble vitamins, and is an important fuel for muscles. Fat is stored in the body in muscle and adipose (fat) tissue in the form of triglycerides which are broken down during exercise into fatty acids that produce energy for muscles. Fat is the primary fuel used in aerobic exercise. Accordingly, one needs fat to burn fat. Fat should be consumed in moderation so the dancer can meet their carbohydrate needs. Excessive fat intake at a given meal will have a negative impact on the dancer's ability to perform fully in class right after the meal as it sits in the stomach for several hours.

Like CHO, however, not all fat from food is the same. Fat can be derived from either animal or plant sources. A cheeseburger with bacon and french fries has a completely different nutrient profile than a meal that includes salmon, brown rice and a salad with avocado. All fats have high calories and can contribute to weight gain if not eaten in moderation; therefore it is important to consider how much fat is eaten on a regular basis. A diet too low in fat can have serious health consequences and ultimately impair performance.

Fats can broadly be divided into saturated and unsaturated based on their chemical structure. While unsaturated fats tend to be found in fish, nuts and seeds, and other plant sources, saturated fats tend to be found in foods of animal origin as well as some manufactured foods. Diets high in some types of saturated fats and/or trans-fats have been shown to contribute to heart disease and cancer.¹³

Processed or fast foods tend to be high in trans-fats and many countries have public health campaigns underway to reduce trans-fatty acid contents in foods. The US FDA has banned trans-fat ingredients from foods. This change will take place within the next few years. Restricting trans-fatty acid intake with continued emphasis on restricting saturated fat intake is recommended.^{14, 15, 16}

While many dancers consume cheeses and full fat dairy products to obtain essential nutrients, these should be consumed in moderation knowing that dairy products also are high in fat. Again, balance is recommended. Foods high in the omega fats play a role in overall cardiovascular health, brain function, and mood. An array of omega-3 and 6 fats are essential to a balanced diet; good sources of omega-3 fats are derived from oily fish, some nuts, linseed/flaxseed oil, and canola oil, whereas omega-6 fats may be found in

vegetable oils. Diets high in the omega fats are beneficial to cholesterol management and heart health. ¹⁷

Research on fat is active and guidance in this area is likely to change. The amount of daily fat needed is approximately 1g per kg body weight, which means that a 50kg female dancer (110 pounds) should eat 45-50g fat over the course of a day, while a 70kg male dancer (154 pounds) should consume around 65-70g fat each day. However, if the dietary goal is weight loss then amounts may need to be lower.

TABLE 2: SOURCES OF FATS

TABLE 2: SOURCES OF FATS				
Food	Amount	Fat content (g)	Notes	Recommendations
Oil, e.g. olive oil	1 level tsp. or 5ml	5g	Mainly monounsaturated.	Use regularly.
Nuts and seeds	1 tbsp. or 15g	10g	Unsaturated and also supply vitamins and minerals.	Limit portion size.
Oily fish, e.g. Salmon (grilled)	Average serving approx. 100g	13g	Mainly unsaturated.	Include 2 portions per week if possible.
Avocado (1/2 large)	100g	8g	Mainly unsaturated.	Treat as a fat source if including regularly.
Butter	1 level tsp. or 5ml	4g	More saturated than unsaturated.	Keep to small amounts.
Nut butter	1 small spoon or 10ml	5g	Mainly unsaturated, fair source of protein.	Watch portion sizes.
Milk – whole	100ml	4g	Mainly saturated – but may not have the health risks of other saturated fats.	Use semi-skimmed for best balance of fats and CHO.
Milk- semi-skimmed/1-2% fat	100ml	1-2g	Lower amount as total fat is lower. Good source of protein,	Include 2-3 servings of milk/yogurt daily.

			calcium and other nutrients.	
Cheddar (hard) cheese	1 slice or 25g		Mainly saturated but good source of calcium and protein.	Make sure a variety of protein rich foods are used.
Mayonnaise	1 tbsp. or 15g	11g	Mainly unsaturated, depending on oil used.	Limit portion size and check amounts in ready-made sandwiches.
Salad dressing	1 tbsp. or 15ml	0-11g	Many brands/recipes – some are fat-free but may have sugar in instead of oil.	Use in small amounts.
Thick/double cream	1 tbsp. or 15g	8g	Mainly saturated.	For occasional use.
Salami	1 slice or 10g	4g	Mixture of fats, high fat, also high salt and preservatives but good source of protein.	For occasional use.
Potato crisps (UK)/chips (USA)	Small bag or 25g	9g	Mainly unsaturated.	Not part of a performance eating plan.
Cake	1 slice or 60g	6-12g approx	Mixture of fats; contains some good nutrients but usually high in calories.	For occasional use.
Biscuits/cookies	2 bourbon cream (24g) 2 packaged chocolate chip cookies	6g	Mixture of fats, energy dense and easy to overeat.	Keep for occasional use.
Chocolate	Small bar approx. 50g	15g	Mainly saturated fat. Limit to high cocoa (dark chocolate) content which has more minerals.	Limit portion size.
Pastry (in pies)	50g	approx . 16g	Varied fat content, high fat to CHO ratio.	Keep for occasional use.

PROTEIN

Protein is a macronutrient and is composed of amino acids, some essential (cannot be made in the human body) and other non-essential (can be made in the body). Amino acids are responsible for the growth of every component and maintenance of every basic function in the human body. Amino acids are used as supplemental fuel to CHO and F (especially when energy supply is insufficient). From the dancer's perspective the importance of protein is its role in repairing muscle fibers that are stressed by constant use in dancing and related activities. Protein is also necessary for bone health. Protein needs are based on body weight rather than on energy requirements for activity. Essential amino acids have to be supplied by food sources which can be derived from either animal or plant sources. Animal proteins provide the most complete array of amino acids, have a higher satiating effect and are more filling because they take longer to digest.¹⁰ Diets higher in protein preserve lean body mass during weight loss.¹⁸ Dancers and athletes are sometimes under the impression that consuming protein powders as a supplement will give them a performance edge or serve as a meal replacement that could be better than food. There is no magic to protein supplements – they deliver good quality protein – nothing more or less. If considering the addition of commercial protein powders, dancers are advised to understand why they are taking them and to verify that the supplements actually contain the ingredients as advertised on the label. Information on purity, safety and contamination guidelines for protein powders can be found online at www.informed-sport.com.




Photo credit: Jasmine Challis

TABLE 3: SOURCES OF PROTEIN

TABLE 3: SOURCES OF PROTEIN		
Animal proteins	Portion size	Protein amount
Chicken, turkey, pork, fish (salmon, tuna)	100g cooked (fish lower than meat) Typical portion size is 100g-150g	20g-30g (fish lower than meat)
Eggs	2 eggs 3 eggs	12g-14g 18g-21g
Cottage cheese, ricotta cheese	100g	10g-12g
Yogurt, Greek yogurt	125g (Greek yogurt has the highest)	6g-10g
Hard cheese	50g	12g
Milk – including whey and casein	1 cup/230ml glass	8g
Plant based proteins	Portion size	Protein amount
Quorn ® (Contains small amount of egg)	100-150g	14g
Split peas, lentils, beans including garbanzo, kidney and butter beans, chickpeas, soy beans, edamame beans	200g ready to use	12g
Wheat gluten/seitan	Seitan 100g	16g
Quinoa	Quinoa raw weight 100g	14g
Soy flour	Soy flour 50g	18g-20g
Tofu	200g	16g
Nuts, seeds and nut butters. These are high in fat; therefore, vary protein sources of protein within a food plan	50g Monitor consumption - check portion size	7g-12g
Soy or Soya milk	1 cup/230ml	5g-9g
'Milk' from hemp, rice, almond, coconut etc., other than soy are much lower in protein, typically 0.1g/100ml therefore, not a useful source of protein	1 cup/230ml glass	Less than 1g

TIMING AND DIGESTION

Scheduling food intake (eating) is almost as important as the type of food and amount that is consumed. It is important to factor in time for food digestion. It is difficult to jump and turn comfortably with a full stomach of steak and eggs, for example, no matter how nutritious the meal. Ideally an interval of 2-4 hours after eating is optimum to allow digestion to take place before dancing. However, most dancers will have to cope with dancing while food is digesting because in the digestive process food takes at least an hour before it leaves the stomach and moves into the small intestine where absorption occurs. It is also nearly impossible to be adequately sustained for an entire day (with an afternoon technique class and evening rehearsal) on an unbalanced diet, whether too little or too much of any given nutrient. Therefore, dancers should research how to consume the type of foods and amounts that will not decrease but increase the physical ability to perform well. The skill is to learn how to balance timing and energy requirement.

Planning tips

Food preparation: Allocate time for grocery shopping, food storage, cooking, and preparation.

Food safety: Be mindful of food temperature; use cool packs if refrigeration is not an option.

Performance and fueling: Maximize energy pre-performance and recovery post performance.

Vacation/Travel: Adjust nutritional requirements when you are not dancing, based on activity.

Injury/post injury: Adjust energy intake to match output and examine balance of nutrients to facilitate healing.

FOOD VS. VITAMINS AND SUPPLEMENTS

Dancers are advised to review their food plans and try to meet vitamin and mineral needs from food. For those with restricted diets or working where food availability is limited, a daily multivitamin and mineral supplement ensures basic requirements are met. However, the addition of vitamin supplements at high dosage to increase performance is not necessary, may impair recovery processes and can be toxic. The exception to this is vitamin D which is covered in greater detail below. To obtain all important macro and micronutrients, a balanced diet composed of a wide variety of fresh fruit and vegetables, whole grains, dairy products, and proteins is recommended. Diets that are restrictive or unbalanced can lead to a range of negative health consequences. Because most individuals do not consistently eat a wide variety of foods, multivitamins are reasonable supplements to take but should not be considered food replacements.

VITAMINS AND MINERALS: MICRONUTRIENTS

Vitamins and minerals comprise the micronutrients in the diet from a wide variety of food and all play a key role in maintaining every system and organ in the body.

Overconsumption can be equally as dangerous as deficiency. Minerals are classified into macro minerals and micro minerals (trace minerals). Although there is this division from a practical point of view they may be considered together, as intakes of all of them are at most a few grams per day (sodium and chloride), and other than calcium (around 1g per day), well under 1 g per day. Many minerals are found in the body but only about 15 are currently known to be essential in our diet, although ongoing research may change this official position in the future. Iron and calcium will be discussed in more detail because of the importance of these minerals.

TABLE 4: MINERALS: SOURCES AND FUNCTION

TABLE 4 MINERALS: SOURCES AND FUNCTION

Minerals most relevant to dance	Functional use	Common source	Potential risk when consumed in extreme
Sodium	Maintains normal blood pressure and water balance, hormonal and muscular function. Usually combined with chloride as salt – (NaCl), generally need to limit as easy to meet requirements.	Table salt, baking soda, seasonings, canned, smoked and salted meats (including bacon and sausages) and fish, olives and pickled foods + processed food (hidden salt)	Both excess and lack of are possible: excess results in thirst (short term); deficiency results in feeling unwell.
Potassium	Maintains normal blood pressure and water balance, muscle function.	Fruit and vegetables, cereals, meat, milk, chocolate, coffee, nuts	Overdose unlikely unless potassium-rich salt is used. Suboptimal intake is common but not immediately harmful.
Chloride	Maintains stomach acidity and fluid balance.	Table salt, soy sauce; large amounts in processed foods; small amounts in milk, meats, breads, and vegetables	Usually combined with sodium, therefore, similar consequences as seen with sodium.
Calcium	Maintains bone health, tooth structure, nerve conduction, and blood clotting.	Milk, yogurt, cheese, fortified soy products, nuts and seeds, green vegetables, dried fruit	Calcium absorption works in conjunction with vitamin D. Absorption is greater when intake is lower, but bone health may be compromised. Blood calcium may rise with very high intakes.
Phosphorous	Maintains bone/tooth formation and energy metabolism.	Milk, yogurt and cheese, grains and cereals, green	Excess/lack is unusual unless caused by physical illness or use with

		vegetables, and meat	certain medications.
Magnesium	Affects muscle contraction, nerve transmission, energy metabolism, and bone integrity.	Nuts and seeds; legumes; leafy, green vegetables; seafood, chocolate; artichokes	Dietary lack/excess unlikely with normal diet.
Iron	Maintains blood integrity; hemoglobin/myoglobin formation to transport oxygen in the body. Ensures a healthy immune system.	Red meat, eggs, cereals, green vegetables, pulses, legumes, dried fruit	Iron deficiency is more common in women. Excess unlikely except when used with certain dietary supplements.
Copper	Involved in enzyme synthesis and varied metabolic functions	Shellfish, nuts, meat including offal, pulses, legumes, cocoa	Dietary lack/excess unlikely with a normal diet.
Selenium	Involved as an anti-oxidant and mediator in electron transfer function.	Brazil nuts, meat, fish, seeds, whole grains	Inadequate amounts are possible; excess amounts are unlikely unless supplements are used. Lack of selenium may trigger mood swings and feelings of depression.
Zinc	Prevents low mood; allows for normal wound healing, and maintains immune system function.	Meat, seafood, green vegetables, seeds	A lack is possible if diet is poor. May cause low mood, poor wound healing, and suppressed immune system. Excess only likely from unwise supplement use.
Manganese	Enzyme synthesis, enzymatic and metabolic functions.	Whole grains, nuts, vegetables, dried fruit, cereals, tea	Dietary lack/excess unlikely with normal diet.

Fluoride	Tooth structure	Seafood, water, tea	Excess possible if use of toothpaste is high. Deficiency unlikely.
Iodine	Thyroid function	Seafood, eggs, dairy	Deficiency is possible depending on soil levels where fruit and vegetables are grown. Excess results from supplement use only.
Chromium	Plays a role in glucose/insulin metabolism.	Whole grains, legumes, lentils, nuts, meat, dairy, eggs	Dietary lack/excess unlikely with normal diet.
Cobalt	Needed as part of Vitamin B12.	Fish, nuts, green leafy vegetables, cereals	Dietary lack/excess is unlikely with normal diet.

CALCIUM

Calcium is important in bone formation. Good bone health is vital because it is important to lay the building blocks for bone stability later in life. During the first two and a half decades of life, bone mass is developed but thereafter, bone formation ceases.^{6,19} It is essential to ingest adequate calcium throughout one's life in addition to the many other nutrients that are required to maintain good bone health. Low bone mass and low calcium intakes are also associated with increased risk of stress fractures. The richest source of calcium is dairy products but generally, can be easily acquired in fortified foods and beverages. Calcium absorption works in conjunction with vitamin D.

IRON

Iron is an essential trace mineral and combined with hemoglobin in red blood cells increases the oxygen capacity of blood. Iron plays an integral role to oxygen storage and transport within the muscle cells. When iron stores are low or deficient, they can be replenished by active iron compounds that are held in reserve in the liver, spleen, and bone

marrow. In the presence of iron insufficiency (anemia), general fatigue, loss of appetite, and inability to perform mild exercise can occur. Iron is absorbed in the intestine; iron derived from animal sources is better absorbed than that from plant sources. Dancers should include normal amounts of iron-rich foods in their daily diet. Dancers who have heavy menstrual periods may need iron supplementation; this need can be confirmed with routine blood work.¹⁹ Dancers who do not eat meat should take care to consume iron from other sources including tuna, egg, oatmeal, dark green leafy vegetables, soy, beans, iron-fortified breakfast cereal, and dried fruits. In the presence of a balanced diet, iron supplements are not needed for most dancers though may be advised for some women.

VITAMINS

Vitamins are organic substances necessary to life and are divided into water soluble and fat soluble. Water soluble vitamins are the B vitamins and vitamin C. Vitamins A, D, E, and K are fat soluble and can be stored in the body which means that they are not required on a daily basis but should be consumed regularly. The fat-soluble vitamin which dancers need to focus on is Vitamin D (See Table 4). Water soluble vitamins, which are Vitamin C and the B group, are needed daily. The B vitamins play important roles in energy production (especially thiamin B1-riboflavin B2-niacin, B3 and B6 – pyridoxine) and in red blood cell formation (folic acid and vitamin B12-cobalamin group). Deficiency of these vitamins can impair performance.

Different fruits and vegetables contain different plant chemicals that can optimize performance as well as serve as anti-oxidants. An easy way to think of this is that different colors in fruit and vegetables represent different effects, so the dancer is well advised to embrace the concept of 'eating across the rainbow'. In general the orange, red, and dark green colored fruits and vegetables supply the highest content of the vitamins A and C. Vitamins A (beta carotene), C, and E function as antioxidants that may help prevent cell damage during exercise and are necessary for the immune system. Smokers are advised to focus on foods, not vitamin supplements for foods rich in anti-oxidants. While

more research is required, there are data that demonstrate that cancer risk in smokers is increased by vitamin supplements (but not by food).²⁰

Vitamin content of food:

<http://ods.od.nih.gov/factsheets/list-VitaminsMinerals/>

Eating across the rainbow:

<http://www.medbroadcast.com/pdf/cccPDF.pdf>

<http://www.ag.ndsu.edu/pubs/yf/foods/fn595.pdf>

VITAMIN D

Many dancers are vitamin D deficient and considered an at-risk population due in part to inadequate or limited sun exposure, increased use of sunblock, and poor diet.²¹ Literature has substantiated that some dancers are deficient in Vitamin D, especially during winter months.²² This deficiency reduces the ability to regenerate muscle and bone function following stress or injury, may interfere with weight loss, and can contribute to the development of stress fractures. Vitamin D recommendations vary largely by country but 400-2000 IU is a safe recommendation for dancers. The UK recommends between 400-1000 IU/ 10-25 mcg, whereas the American College of Sports Medicine recommends 1000-2000 IU during the winter months, and the Australian recommendation is 600-4000 IU. Recent evidence suggests that approximately 1000 IU are needed to maintain adequate levels of vitamin D.²² A safe upper limit seems to be 4000 IU/100micrograms for the USA and Europe and 3200 IU/80 micrograms for Australia. A study in which dancers were given 2000 IU Vitamin D for 4 months yielded beneficial results. The authors also noted however, that if blood vitamin D levels are not monitored, 1000 IU per day is recommended.²³ This fits within a safe upper limit of 4000 IU/100micrograms (3200 IU US and Europe recommendations and 800 micrograms from Australia). Vitamin D supplementation has also been associated with increased vertical jump height and isometric strength, and lower injury rates among elite ballet dancers.²¹

It is recommended that dancers spend some time outside (at least 10 minutes per day), with no hat, sunscreen, and sleeves rolled up to optimize Vitamin D exposure. In addition, dancers are encouraged to seek the advice of their local medical professional regarding the use of Vitamin D supplementation.

See also the IADMS resource paper on bone health <http://www.iadms.org/?212>.

For more information on supplements around the world

<https://www.fda.gov/food/dietarysupplements/>

http://www.nhs.uk/news/2011/05may/documents/BtH_supplements.pdf

<http://www.nutritionaustralia.org/national/resources/sports-nutrition>

<http://www.efsa.europa.eu/en/topics/topic/drv.htm>

SPICES

Spices and herbs have been used for centuries in medicine and health. Bioactive compounds in herbs and spices, collectively called, dietary phenols or phytonutrients, contain properties that have been purported to reduce inflammation and oxidative stress, improve cardiovascular and metabolic function, cognition, gut microbiota, digestion, and reduce risk of certain cancers. Polyphenols and polyphenol rich foods especially fruits, vegetables, and green tea are known for their antioxidant properties. In normal amounts they may have more of a role in overall health and maintenance than in the development or prevention of disease (e.g. cinnamon, turmeric, etc.). In larger amounts they may interact with medications and may have a negative effect on physiological function.^{24, 25, 26}

COMPLEMENTARY AND ALTERNATIVE MEDICINE

Complementary, homeopathic, and alternative medicine treatments should be approached with caution, and reviewed by credible sources before embarking on potentially expensive regimens. Herbs (e.g. St. John's Wort) and media-hyped cleanses and teas are not

recommended as routine practice as they can interfere with medications and create unwanted or harmful effects even in the absence of medications. Medicinal or herbal teas can have either diuretic or laxative properties and should not be used to modify weight.

3. FLUID, HYDRATION, AND SWEAT

When we dance or exercise, heat is generated by muscles and raises core temperature. Within normal limits, increased core temperature as a result of dancing does not lead to impaired thermal (heat) regulation. Exercise increases heat production in the muscles and cooling the body is primarily dependent on the evaporation of sweat from the skin. Perspiration is a normal body function and allows the body to regulate its ability to adapt to exercise and hydration. The amount of sweat produced may vary by person, and a 16-50 ounce (0.5-1.5-liter) loss during moderate exercise over a one hour period is common. With back to back classes, dancers can lose a considerable amount of water through sweat.

Water accounts for 60% of the total weight of the human body. Dancers need to stay hydrated; without proper hydration fatigue and injury can result. Environmental temperature and sweat production will drive the amount of fluid needed to maintain health. Sweat losses during training or performance from exertion can be substantial and vary by sport and gender. Dehydration of 3% of body weight can lead to cramps, nausea, light headedness or fainting and may severely impair performance. Adequate fluid replacement prevents dehydration and its consequences.²⁷ A well-hydrated body will produce a good volume of urine that is pale in color and does not have a strong odor. As a self-test, drink 400-600 ml two hours before dance and check urine output over the next hour.²⁸ Dancers should note that B vitamin supplements may produce a stronger yellow color which should be considered in evaluation.²⁹

Since many dance environments are kept warm because dancers prefer warm temperatures to keep their muscles malleable and often add layers of warm (breathable) clothing, sweating has an additive effect on fluid loss. Therefore, it is vital to replace fluids

often. The amount of fluid loss is related to body size, mass, and genetic factors, with highest losses from taller heavier males, and lowest losses in lighter females.³⁰

HYDRATION TIPS

- Stay hydrated by drinking non-sugared liquids. Don't restrict fluids during meals or exercise.
- Drink frequently throughout the day 'ad lib' or 'to thirst', even if not thirsty - keep a full, non-spill bottle at hand.
- Drink more to delay hydration and diminish a rise in core temperature.
- Water is the best replacement for sweat loss as sweat is mostly water and sodium.
- Water is easy to obtain, has no additives, and is quickly absorbed.
- Sodium replacement is not necessary, but for those who sweat profusely, electrolytes may be recommended. Salt restricted diets should only be advised for medical reasons.

ELECTROLYTES

Electrolytes are minerals the body requires for regulating water balance, blood acidity, and muscle function. They act as conductors within body fluid that have both positive and negative charges that keep the body fluid in balance and play a role in every physiologic system. The major electrolytes are located in extracellular fluid and include sodium, potassium, calcium, and magnesium. These in turn are balanced by a host of concentrations that reside primarily in the kidney.³¹ Changes in food intake (fasting, re-feeding, limiting calories, etc.) severely impacts water and electrolyte balance which can elevate thirst, body weight, and water retention - conditions dancers do not want to have fluctuating arbitrarily.

The major source of sodium in the diet is added salt. One teaspoon of salt is equal to 2,300 mg (2.3 g) of sodium. Because salt is so widely used in our food today, inattention to the amount of sodium in one's diet can easily lead an imbalance which can impact not only how the body looks and feels, but how it performs as well.

COMMON BEVERAGES

Teas and Coffee

Coffees and teas are best consumed in moderation as caffeine is a stimulant. Moderate intake of caffeine has been shown to enhance endurance and sports performance but can also cause jitteriness when consumed in excess. While caffeine has mild diuretic properties, this response is seen in people who do not regularly consume caffeine and the effect is moderated with continued consumption.³² As caffeine lingers in the body for several hours it is best to limit consumption to the morning or afternoon and limit in the early evening in order to obtain a good night's sleep. Caffeinated drinks should not be used to control appetite. Repackaged and commercial teas and coffees often include unwanted high levels of sugar, sodium, calories, and artificial ingredients so always check the ingredients.

Fruit drinks and juices

In the US, fruit juices may be fortified with vitamins and nutrients. In general, juices with pulp are more nutritious compared to those without. On the other hand, fruit drinks can be sources of high sugar and additives and have variable amounts of fruit juice in them. Drinking large amounts of juice at meals and throughout the day should be avoided as it can contribute to weight gain, may not satisfy thirst, and may make it harder to regulate appetite. The acidity and sugar content associated with juice can have a deleterious effect on oral health.³³

Soda

Soda and other carbonated drinks can be high in sugar and high fructose corn syrup, artificial additives, sodium, caffeine, and contribute to tooth decay and interfere with digestion. While drinks and sodas made with artificial sweeteners, collectively called non-

nutritive sweeteners, may have zero calories, it is a subject of debate within the scientific community whether they impact the brain to perceive them as metabolically active and interfere with weight regulation.^{34, 35} There are no health benefits to soda and they are best avoided before and during dance as the carbonation can cause stomach discomfort and gas.

Energy & sport drinks

'Energy' drinks are high in caffeine, artificial additives, and sugar and offer very little nutritional value and are therefore, not recommended for dancers. The extreme amounts of sugar (3 teaspoons per 4 ounce or 15g per 100ml) and caffeine in these drinks can cause excessive jitteriness, loss of focus and concentration. They can also trigger previously unrecognized heart conditions, headache or migraine and are particularly dangerous when mixed with alcohol as alcohol is depressant to the nervous system, and mixed with caffeine, a stimulant. For more information, follow this link:

<https://www.sportsdietitians.com.au/factsheets/children/nutrition-for-the-adolescent-athlete/>

Commercial sports drinks are often promoted as good fluid replacements during high intensity, prolonged activities in which water and electrolytes may be radically depleted, and where additional carbohydrates may be required. Sports drinks can be useful to dancers with high fuel (caloric) needs (some male dancers and smaller number of female dancers), those who perspire heavily, have a diet naturally low in salt or train in very hot environments. These drinks are isotonic (contain a similar number of salts and carbohydrates particles as bodily fluids) and supply approximately one teaspoon of CHO as sugar per 4 ounce (4-6g per 100ml), plus salt, which translates to 16-24kcal per 4 ounce/ 100ml. The effect of vitamin loss in sweat in dancing is likely to be negligible.³⁶ As a general rule, commercial sports drinks are not needed in dance.

Protein shakes and smoothies

While American audiences mostly recognize shakes as made primarily with ice cream and syrup, this section discusses shakes in the context of those prepared with healthy additives like protein, antioxidants, or vitamins and made from water, juice, or milk. Smoothies and protein shakes are frequently marketed as nutritious and used as meal replacements but can vary widely in ingredients. Smoothies and shakes should be considered as part of a meal and not as a meal replacement. Although a smoothie can provide several pieces of fruit, it is easy to over-consume, because it is difficult to ascertain how much fruit is in a typical serving. Even the natural sugar in fruit can be too much when consumed in excess. A healthier alternative is to make one's own smoothie and include raw vegetables. The blended fiber can be a useful part of a meal plan as well and the addition of milk/soy milk/yogurt can make a useful recovery drink. With that in mind, drink immediately or soon after preparing to avoid vitamin loss which ultimately lowers the potential health benefits.

Supplement-based shakes also vary widely in their content and thus their potential benefit. They can contain high amounts of fat and sugars and contribute little to an eating plan. In contrast, shakes made with low fat milk or a milk substitute (soy, cashew, almond, rice, hemp, etc.) and combined with added fruit and protein powder or dried skimmed milk powder (if needed) are a better alternative that have more healthy ingredients and less added sugar. Commercial milkshakes like smoothies or iced coffee drinks vary in both the size they are sold in and their composition. Drinks made with high sugar ingredients are not as nutrient-rich, compared to those made primarily from milk and fruit. Consider also that frozen fruit is a great alternative and less costly when fresh fruit is unavailable. Always check the ingredients' labels and limit the size.

HOME- MADE SPORTS/ISOTONIC DRINK RECIPES*

Variations can be found online. Keep the ratios of ingredients in similar proportion and modify to taste.

Isotonic drink based on juice concentrate, fruit squash or cordial (fruit flavored concentrates either with sugar or sweetener) reconstituted in a ratio of around 1 part concentrate to 6 parts water.

7 oz. (200ml) ordinary juice concentrate

28 oz. (3 ½ cups) or 800ml water

Pinch of salt

Mix together in a large container and refrigerate.

Makes ~5 cups

Isotonic drink based on pure/100% fruit juice

Approximately 16 ounces (500ml) unsweetened fruit juice (orange, apple, pineapple)

Approximately 16 ounces (500ml) water

Mix together in a large container and refrigerate.

Makes 4 cups

Isotonic drink based on glucose/sugar plus flavour according to taste

3-4 tbsps. (50-70g) sugar or glucose powder

32 ounces (1 litre) warm water

Pinch of salt

Up to 7 ounces or 200ml of sugar free concentrate or 1 drink flavour packet according to taste

Mix and refrigerate

Makes 4 cups

* Isotonic drinks have comparable salts and carbohydrates to body composition

Alcohol

Alcoholic drinks do not enhance performance and have dehydrating properties that counteract dancers' efforts to balance fluids. Alcohol is a toxic substance and provides no vitamins, minerals or proteins and moderate consumption may be a cancer risk. Even small amounts of alcohol (15 grams or 6 oz.) can increase one's cancer risk and amounts greater than this offset any potential benefits.³⁷ Mixers added to alcoholic drinks are mostly CHO in the form of sugar which is rapidly absorbed into the bloodstream. When a carbonated mixer is added, absorption rate may vary, leaving the dancer unexpectedly vulnerable to toxicity and in extreme cases, alcohol poisoning.

Chart I: Pre/post performance meal planning

Chart 1: PRE/POST PERFORMANCE MEAL PLANNING

	During long training days (rehearsals, multiple classes with few breaks between)	1-2 hours prior to dancing (short term energy needed)	Following high exertion (within 1.5 hours post performance)
Rationale	<ul style="list-style-type: none"> • Maintain energy • Prevent fatigue • Maintain satiety/minimize hunger 	<ul style="list-style-type: none"> • Provide quick energy • Digest quickly and empties stomach quickly. 	<ul style="list-style-type: none"> • Replenish muscle glycogen and muscle synthesis. • Provide energy for the next day's activity.
Food examples	<ul style="list-style-type: none"> • Items for meals and snacks: e.g. couscous/quinoa • Bread with protein plus vegetables, fruit/ dried fruit with nuts/seeds • Cereal or protein bar, handful of wholegrain cereal 	Fruit e.g. banana with a few nuts/seeds; oat cakes with hummus or cereal bar or chicken/beef/egg sandwich or natural yogurt topped with nuts and fruit	<ul style="list-style-type: none"> • Milk-based or soy drink/shake or yogurt with fruit. • Follow up with meal containing CHO, P and some F.

Chart 2: Sample food plan for one day

Chart 2: SAMPLE FOOD PLAN FOR ONE DAY

Meal or snack	Suggestions	Timing	Rationale
Breakfast	Oat cereal with milk, fruit and nuts/seeds	At least 1 hour but ideally 2 hours before first class.	Ensures energy supplies are going to be steady through the morning.
Break	Fruit – adding nuts for more strenuous days	Any short break if a long training day.	Tops off energy levels especially if the day is strenuous.
Lunch	Some protein balanced with carbohydrates and fats, e.g. veg/salad, avocado/nuts	As early as possible in the lunch break to allow maximum digestion before	Replenishes nutrients used in the morning and fuel for afternoon/evening activities.

		the next class/rehearsal.	
Break	Fruit or nuts	Any short break if a long training day.	Tops off energy levels, especially if the day is strenuous.
Dinner	Some protein balanced with carbohydrates and fat, e.g. veg/salad, avocado/nuts	As soon as possible after last class/rehearsal.	Facilitates ongoing replenishment and muscle repair.
Break	Milky drink/bowl of cereal/herbal tea	Late evening/before bed.	Allows those with higher requirements to choose appropriate food and others a calming drink before sleep.

RECOMMENDATIONS: TAKE HOME MESSAGES

- Fuel appropriately for all activities: eat breakfast soon after waking up and prior to training.
 - Seek out a variety of foods and fuel sources for optimum performance nutrition.
 - Diets do not work. Healthy eating is an enjoyable lifelong process.
 - Always check the ingredients' labels and monitor portions.
 - Drink water frequently and stay hydrated with meals.
 - Be prepared: plan healthy meals and snacks in advance to ensure optimum energy levels.
 - Wherever possible, eat foods in their natural form rather than highly processed alternatives.
 - Prioritize sleep quality and quantity; balance rest and daily training.
- 80% of injury recovery is due to adequate rest, nutrition and sleep.* ³⁸
- Talk to a teacher or parent if eating behavior feels out of control.

4. RECOMMENDED READING

Nutrition Books – sports/dance - based:

All of the books listed below provide great information for dancers. This list is not exhaustive but aims to provide a global selection of resources. As different books, styles and balance of theory to practice suit different individuals, there is no “best” book; it will be down to individual preference to adapt the information to one’s own practice.

Albers S. *Eating Mindfully*, 2nd ed. California: New Harbinger Publications, 2012.

Bean A. *Food for Fitness: How to Eat for Maximum Performance*, 4th ed. London: Bloomsbury Sport, 2014.

Bean A. *The Complete Guide to Sports Nutrition*, 7th ed. London: Bloomsbury Sport, 2013.

Burke L. *The Complete Guide to Food for Sports Performance*, 3rd ed. Australia: Allen & Unwin, 2010.

Burke L, Deakin V. *Clinical Sports Nutrition*, 5th ed. Australia: McGraw-Hill Education, 2015.

Cardwell G. *Gold Medal Nutrition*, 5th ed. Champaign, IL: Human Kinetics, 2012.

Clark N. *Nancy Clark’s Sports Nutrition Guidebook*, 5th ed. Champaign, IL, 2013.

Costa R. *Cooking for Sport and Exercise*. UK: Coventry University Enterprises, 2012.

Gibney MJ, Lanham-New SA, Cassidy A, Vorster, HH (eds.) *Introduction to human nutrition*. Oxford: John Wiley & Sons, 2013.

Jukendrop A, Gleesen M. *Sports Nutrition*, 2nd ed. Champaign, IL: Human Kinetics, 2010.

Lanham-New S, Stear S, Shirreffs S, Collins A. *Sport and Exercise Nutrition*. Oxford: Wiley-Blackwell, 2011.

Mastin Z. *Nutrition for the Dancer*. London: Dance Books Ltd, 2009.

5. REFERENCES

1. Arcelus J, Witcomb GL, Mitchell A. Prevalence of Eating Disorders amongst Dancers: A Systemic Review and Meta-Analysis. *Eur. Eat. Disorders Rev.* 2014;22: 92–101. doi: 10.1002/erv.2271.
2. Brown D & Wyon M. An International Study on Dietary Supplementation Use in Dancers. *Med Probl Perform Art.* 2014;29(4): 229–234.
3. Hu FB, Satija A, Manson JE. Curbing the Diabetes Pandemic: The Need for Global Policy Solutions. *JAMA.* 2015;313(23):2319-2320.
4. Eating Disorders, National Institute of Health. Available at: <http://www.nimh.nih.gov/health/topics/eating-disorders/index.shtml>.
5. The Prix de Lausanne Health Policy. Available at: <http://www.prixdelausanne.org/competition/health-policy/>
6. Manore M, Meyer NL, Thompson J. *Sport nutrition for health and performance.* Champaign, IL: Human Kinetics, 2009.
7. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain DP. Quantity and Quality of Exercise for Developing and Maintaining Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Apparently Healthy Adults: Guidance for Prescribing Exercise. 2011 ACSM Position Stand. *Med Sci Sports Exerc.* 2011;43(7):1334-1359. doi: 10.1249/MSS.0b013e318213febf.
8. Desbrow B, McCormack J, Burke LM, Cox GR, Fallon K, Hislop M, Logan R, Marino N, Sawyer SM, Shaw, G, Star A, Vidgen H, Leveritt M. Sports Dietitians Australia position statement: sports nutrition for the adolescent athlete. *Int J Sport Nutr Exerc Metab.* 2014;24(5):570-584. doi: 10.1123/ijsnem.2014-0031.
9. De Souza MJ, Nattiv A, Joy E, Misra M, Williams NI, Mallinson RJ, Gibbs JC, Olmsted M, Goolsby M, Matheson G. 2014 Female Athlete Triad Coalition Consensus Statement on Treatment and Return to Play of the Female Athlete Triad: 1st International Conference held in San Francisco, California, May 2012 and 2nd International Conference held in Indianapolis, Indiana, May 2013. *Br J Sports Med.* 2014;48:289. doi:10.1136/bjsports-2013-093218
10. Mountjoy M, Sundgot-Borgen JK, Burke LM, *et al* IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Br J Sports Med* 2018; 52:687-697
11. Westerterp-Plantenga MS, Lemmens SG and Westerterp KR. Dietary protein – its role in satiety, energetics, weight loss and healthy eating. *Br J Nutrition.* 2012;108, S105–S112. doi:10.1017/S000

12. Baker LB, Rollo I, Stein KW, Jeukendrup AE. Acute Effects of Carbohydrate Supplementation on Intermittent Sports Performance. *Nutrients*. 2015;7, 5733-5763; doi:[10.3390/nu7075249](https://doi.org/10.3390/nu7075249)
13. Hooper L, Martin N, Abdelhamid A, Davey Smith G. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database Syst Rev*. 2015;10:6. doi: 0.1002/14651858.CD011737.
14. Dawczynski C, Kleber ME, März W, Jahreis G, Lorkowski S. Saturated fatty acids are not off the hook. *Nutr Metab Cardiovasc Dis*. 2015; Oct 9:S0939-4753(15)00218-5. doi: 10.1016/j.numecd.2015.09.010.
15. Lichtenstein A. Dietary trans fatty acids and cardiovascular disease risk: past and present. *Curr Atheroscler Rep*. 2014;16(8):433. doi: 10.1007/s11883-014-0433.
16. FDA Cuts Trans Fat in Processed foods: Federal Drug Administration: <http://www.fda.gov/ForConsumers/ConsumerUpdates/ucm372915.htm>
17. Lorente-Cebrian S, Costa AG, Navas-Carretero S, Zabala M, Martinez JA, Moreno-Aliaga MJ. Role of Omega-3 fatty acids in obesity, metabolic syndrome, and cardiovascular diseases: A review of the evidence. *J Physio and Biochem*. 2013; 69: 633-651. doi: 10.1007/s13105-013-0265-4
18. Mettler S, Mitchell N, Tipton KD. Increased protein intake reduces lean body mass loss during weight loss in athletes. *Med Sci Sports Ex* 2010;42(2):326-337 doi: 10.1249/MSS.0b013e3181b2ef8e.
19. Burke L. *Practical sports nutrition*. Champaign, IL: Human Kinetics, 2007
20. Harvie M. Nutritional supplements and cancer: potential benefits and proven harms. *Am Soc Clin Oncol Educ Book*. 2014:e478-486. doi: 10.14694/EdBook_AM.2014.34.e478.
21. Wyon MA, Koutedakis Y, Wolman R, Nevil AM, Allen N. The Influence of winter vitamin D supplements on muscle function and injury occurrence in elite ballet dancers. *J Sci Med Sport* 2014; 17(1):8-12. doi: 10.1016/j.jsams.2013.03.007.
22. Wolman R, Wyon MA, Koutedakis Y, Nevill AM, Eastell R, Allen N. Vitamin D status in professional ballet dancers: winter vs. summer. *J Sci Med Sport*. 2013; 16(5):388-391. doi: 10.1016/j.jsams.2012.12.010.
23. Wolman R. personal communication, 2014.
24. Opara EI, Chohan M. Culinary Herbs and Spices: Their Bioactive Properties, the Contribution of Polyphenols and the Challenges in Deducing Their True Health Benefits. *Int. J. Mol. Sci*. 2014; 15: 19183-19202; doi:10.3390
25. Rubi L, Motilva MJ and Romero MP. Recent Advances in Biologically Active Compounds in Herbs and Spices: A Review of the Most Effective Antioxidant and Anti-Inflammatory Active Principles. *Crit Rev Food Sci Nutr*. 2013;53(9): 943-953 doi: 10.1080/10408398.2011.574802.

26. Morrison J. Mealographer website
http://www.mealographer.com/food.php?action=search&order_1=high&display_name_1=protein&order_2=&display_name_2=Calories&FdGrp_Desc=Spices+and+Herbs&submit=Submit
27. Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. American College of Sports Medicine position stand: Exercise and fluid replacement. *Med Sci Sport Ex.* 2007; 39(2): 377-390.
28. Latzka WA, Montain SJ. Water and electrolyte requirements for exercise. *Clin Sports Med.* 1999;18(3):513-524.
29. Hew-Butler T, Rosner MH, Fowkes-Godek S, Dugas JP, Hoffman MD, Lewis DP, Maughan RJ, Miller KC, Montain SJ, Rehrer NJ, Roberts WO, Rogers IR, Siegel AJ, Stuempfle KJ, Winger JM, Verbalis JG. Statement of the Third International Exercise-Associated Hyponatremia Consensus Development Conference, Carlsbad, California, 2015. *Clin J Sport Med.* 2015;25(4):303-320. doi: 10.1097/JSM.0000000000000221.
30. Maresh CM, Gabaree-Boulant CL, Armstrong LE, Judelson DA, Hoffman JR, Castellani JW, Kenefick RW, Bergeron MF, Casa DJ. Effect of hydration status on thirst, drinking, and related hormonal responses during low-intensity exercise in the heat. *J Appl Physiol.* 2004; 97(1):39-44. DOI: 10.1152/jappphysiol.00956.2003
31. Shirreffs SM, Sawka MN. Fluid and electrolyte needs for training, competition, and recovery. *J Sports Sci.* 2011; 29:sup1, S39-S46, DOI: 10.1080/02640414.2011.614269
32. Maughan RJ, Griffin J. Caffeine ingestion and fluid balance: a review. *J Hum Nutr Diet.* 2003; 6(6):411-420.
33. Dugmore CR and Rock WP. A multifactorial analysis of factors associated with dental erosion *Br Dent J.* 196, 283 - 286 (2004) doi:10.1038/sj.bdj.4811041.
34. Burke MV, Small DM. Physiological mechanisms by which non-nutritive sweeteners may impact body weight and metabolism. *Physiol Behav.* 2015; 152(Pt B):381-388. doi: 10.1016/j.physbeh.2015.05.036.
35. Pepino YM. Metabolic effects of non-nutritive sweeteners. *Physiol Behav.* 2015;152: 450–455.
36. Peng Y, Cui X, Liu Y, Li Y, Jian Liu J, Cheng B. Systematic Review Focusing on the Excretion and Protection Roles of Sweat in the Skin. *Dermatology;* 2014: 228:115–120 DOI: 10.1159/000357524
37. Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, Scott L, Jenab M, Turati F, Pasquali E, Pelucchi C, Bellocco R, Negri E, Corraro G, Rehm J, Boffett P,

- LaVecchia C. Light alcohol drinking and cancer: a meta-analysis. *Ann Oncol.* 2013;24(3):301–308.
38. Rosenblatt B. personal communication, 2015.

6. AUTHORS

Jasmine Challis is an RD, and
Adrienne Stevens, EdD is